




GOAT ANTI-CSF1R (AA734-744) ANTIBODY

SKU: EB10258



SPECIFICATIONS

Formulation Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Unit Size 100 µg

Storage Instructions Aliquot and store at -20°C. Minimize freezing and thawing.

Synonym / Alias Names CSF1R|McDonough feline sarcoma viral (v-fms) oncogene homolog|macrophage colony stimulating factor 1 receptor|FMS proto-oncogene|FMS|FIM2|CSFR|colony stimulating factor 1 receptor|C-FMS|CD115 antigen|CD115

Usage Summary

<p>Flow Cytometry: Flow cytometric analysis of THP-1 cells. Recommended concentration: 10ug/ml.</p>

Accession ID NP_005202.2; NP_001275634.1; NP_001362250.1

Blocking Peptide EBP10258

Immunogen Peptide with sequence C-NDSFSEQDLDK, from the internal region of the protein sequence according to NP_005202.2; NP_001275634.1; NP_001362250.1.

Peptide Sequence C-NDSFSEQDLDK





Purification Method	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Shipping Instructions	Refrigerated
Predicted Species	Human, Dog, Cow
Reactive Species	Human
Human Gene ID	1436
Rat Gene ID	307403
Product Grade	https://prod-vector-labs-pimcore-assets.s3.us-east-1.amazonaws.com/assets/products/image/elite_medium.png
IHC Results	Paraffin embedded Human Spleen and Tonsil. Recommended concentration: 5-6µg/ml.
ELISA Detection Limit	Antibody detection limit dilution 1:8000.
Western Blot	Approx. 100kDa band observed in lysates of cell line THP-1 (calculated MW. of 108kDa according to NP_005202.2). Recommended concentration: 1.5-3µg/ml. Primary incubation 1 hour at room temperature.
Application Type	Pep-ELISA, FC, IHC, WB

DOCUMENTS

- [Data Sheet](#)

GALLERY IMAGES





www.everestbiotech.com

Email: customerservice@vectorlabs.com

Telephone: [\(650\) 697-3600](tel:(650)697-3600)

